

On the two different bi-Hamiltonian structures for the Toda lattice.

A V Tsiganov

St.Petersburg State University, St.Petersburg, Russia

e-mail: tsiganov@mph.phys.spbu.ru

Abstract

We construct two different incompatible Poisson pencils for the Toda lattice by using known variables of separation proposed by Moser and by Sklyanin.

1 Introduction

A bi-Hamiltonian manifold M is a smooth manifold endowed with two compatible bi-vectors P_0, P_1 such that

$$[[P_0, P_0]] = [[P_0, P_1]] = [[P_1, P_1]] = 0,$$

where $[[\cdot, \cdot]]$ is the Schouten bracket. Such a condition assures that the linear combination $P_0 - \lambda P_1$ is a Poisson pencil, i.e. it is a Poisson bi-vector for each $\lambda \in \mathbb{C}$, and therefore the corresponding bracket $\{.,.\}_0 - \lambda\{.,.\}_1$ is a pencil of Poisson brackets [5].

Dynamical systems on M having enough functionally independent integrals of motion $H_1, \dots, H_n$ in involution with respect to the both Poisson brackets

$$\{H_i, H_j\}_0 = \{H_i, H_j\}_1 = 0. \quad (1.1)$$

will be called bi-integrable systems.

The main aim of this note is to prove that any separable system is bi-integrable system. As an example we show that the Toda lattice is bi-integrable system with respect to two different Poisson pencils $P_0 + \lambda P_1$ and $P_0 + \lambda P_1^*$, which are related with two known families of the separated variables [6, 7].

2 The separation of variables method

A complete integral $S(q, t)$ of the Hamilton-Jacobi equation

$$\frac{\partial S(q, t)}{\partial t} + H\left(q, \frac{\partial S(q, t)}{\partial q}, t\right) = 0, \quad (2.1)$$

where $q = (q_1, \dots, q_n)$, is a solution $S(q, t, \alpha_1, \dots, \alpha_n)$ depending on n parameters $\alpha = (\alpha_1, \dots, \alpha_n)$ such that

$$\det \left\| \frac{\partial^2 S(q, t, \alpha)}{\partial q_i \partial \alpha_j} \right\| \neq 0. \quad (2.2)$$

For any complete integral of (2.1) solutions $q_i = q_i(t, \alpha, \beta)$ and $p_i = p_i(t, \alpha, \beta)$ of the Hamilton equations of motion may be found from the Jacobi equations

$$\beta_i = -\frac{\partial S(q, t, \alpha)}{\partial \alpha_i}, \quad p_i = \frac{\partial S(q, t, \alpha)}{\partial q_i}, \quad i = 1, \dots, n. \quad (2.3)$$

According to (2.2) if we resolve a second part of the Jacobi equations with respect to parameters $\alpha_1, \dots, \alpha_n$ one gets n independent integrals of motion

$$\alpha_m = H_m(p, q, t), \quad m = 1, \dots, n, \quad (2.4)$$

as functions on the phase space M with coordinates p, q .

Definition 1 *A dynamical system is a separable system if the corresponding complete integral $S(q, t, \alpha)$ has an additive form*

$$S(q, t, \alpha_1, \dots, \alpha_n) = -Ht + \sum_{i=1}^n S_i(q_i, \alpha_1, \dots, \alpha_n). \quad (2.5)$$

Here the i -th component S_i depends only on the i -th coordinate q_i and α .

In such a case Hamiltonian H is said to be separable and coordinates q are said to be separated coordinates for H , in order to stress that the possibility to find an additive complete integral of (2.5) depends on the choice of the coordinates.

For the separable dynamical system we have

$$\frac{\partial^2}{\partial q_k \partial q_j} S = \frac{\partial}{\partial q_k} \left(\frac{\partial}{\partial q_j} S_j \right) = 0, \quad \text{for all } j \neq k, \quad (2.6)$$

such that the second Jacobi equations (2.3) are the separated equations

$$p_j = \frac{\partial}{\partial q_j} S_j(q_j, \alpha_1, \dots, \alpha_n), \quad \text{or} \quad \phi_j(p_j, q_j, \alpha) = p_j - \frac{\partial}{\partial q_j} S_j(q_j, \alpha) = 0. \quad (2.7)$$

Here j -th equation contains a pair of canonical variables p_j and q_j only.

Proposition 1 *For any separable dynamical system integrals of motion $H_m(p, q, t)$ (2.4) are in the involution*

$$\{H_k, H_m\}_f = 0, \quad k, m = 1, \dots, n,$$

with respect to the following brackets $\{.,.\}_f$ on M

$$\{p_i, q_j\}_f = \delta_{ij} f_j(p, q), \quad \{p_i, p_j\}_f = \{q_i, q_j\}_f = 0, \quad (2.8)$$

which depend on n arbitrary functions $f_1(p, q), \dots, f_n(p, q)$.

PROOF: In fact we have to repeat the proof of the Jacobi theorem due by Liouville. Namely, differentiate the relations (2.4) by q_j and then substitute momenta p_k from separated equations (2.7) to obtain

$$\frac{\partial H_m}{\partial q_j} + \sum_{i=1}^n \frac{\partial H_m}{\partial p_i} \frac{\partial p_i}{\partial q_j} = \left(\frac{\partial H_m}{\partial q_j} + \frac{\partial H_m}{\partial p_j} \frac{\partial^2 S_j}{\partial q_j^2} \right) = 0.$$

It follows that for any H_k and H_m

$$\sum_{j=1}^n f_j \frac{\partial H_k}{\partial p_j} \left(\frac{\partial H_m}{\partial q_j} + \frac{\partial H_m}{\partial p_j} \frac{\partial^2 S_j}{\partial q_j^2} \right) = \sum_{j=1}^n f_j \frac{\partial H_k}{\partial p_j} \frac{\partial H_m}{\partial q_j} + \sum_{j=1}^n f_j \frac{\partial H_k}{\partial p_j} \frac{\partial H_m}{\partial p_j} \frac{\partial^2 S_j}{\partial q_j^2} = 0.$$

Permute indexes k and m and subtract the resulting equation from the previous one we get

$$\sum_{j=1}^n f_j \left(\frac{\partial H_k}{\partial p_j} \frac{\partial H_m}{\partial q_j} - \frac{\partial H_m}{\partial p_j} \frac{\partial H_k}{\partial q_j} \right) = 0.$$

The final assertion easily follows.

Brackets $\{.,.\}_f$ (2.8) are the Poisson brackets if and only if

$$[[P_f, P_f]] = 0, \quad (2.9)$$

where

$$P_f = \begin{pmatrix} 0 & \text{diag}(f_1, \dots, f_n) \\ -\text{diag}(f_1, \dots, f_n) & 0 \end{pmatrix}. \quad (2.10)$$

Proposition 2 *If j -th function*

$$f_j(p, q) = f_j(p_j, q_j), \quad j = 1, \dots, n, \quad (2.11)$$

depends only on the j -th pair of coordinates p_j, q_j then brackets $\{.,.\}_f$ (2.8) are the Poisson brackets, which are compatible with canonical ones.

PROOF: Substitute tensor P_f (2.10) into the equations

$$[[P_0, P_f]] = [[P_f, P_f]] = 0, \quad \text{where} \quad P_0 = \begin{pmatrix} 0 & \mathbf{I} \\ -\mathbf{I} & 0 \end{pmatrix},$$

one gets the following system of partial differential equations

$$\frac{\partial f_j}{\partial q_k} = \frac{\partial f_j}{\partial p_k} = 0, \quad f_i \frac{\partial f_j}{\partial q_k} = f_i \frac{\partial f_j}{\partial p_k} = 0,$$

for all $j \neq k \neq i$. The separable functions $f_j(p_j, q_j)$ (2.11) satisfy this system of equations. So, the corresponding tensor P_f (2.10) is the Poisson tensor, which is compatible with canonical tensor P_0 .

Remark 1 Separated variables (p, q) are defined up to canonical transformations

$$p_j = X(p_j, q_j), \quad q_j = Y(p_j, q_j),$$

which have to preserve canonical tensor P_0 and would change the second tensor P_f . It is clear, that freedom in the choice of the functions $f_j(p_j, q_j)$ is related with this freedom in the definition of the separated variables.

According to the following proposition, the separation of variables method is closely related with the bi-Hamiltonian geometry.

Proposition 3 *Any separable dynamical system is bi-integrable system.*

PROOF: In order to get a pair of compatible Poisson brackets on the phase space M it is enough to postulate brackets (2.8)

$$\{p_i, q_j\}_f = \delta_{ij} f_j(p_j, q_j), \quad \{p_i, p_j\}_f = \{q_i, q_j\}_f = 0, \quad (2.12)$$

between the known separated variables q_j and p_j for a given dynamical system. According to the Proposition 1 the corresponding integrals of motion $H_m(p, q, t)$ (2.4) are in the bi-involution with respect to brackets $\{.,.\}_0$ and $\{.,.\}_f$.

In the next sections we demonstrate how this proposition works for the open and periodic Toda lattice.

Remark 2 The separated variables (p, q) are canonical variables, which put the corresponding recursion operator $N = P_f P_0^{-1}$ in diagonal form. So, according to [2, 3], a set of the separated variables (p, q) is said to be Darboux-Nijenhuis coordinates.

Remark 3 For the stationary systems differentiate the separated equations (2.7)

$$\left(\frac{\partial \phi_j}{\partial q_j} dq_j + \frac{\partial \phi_j}{\partial p_j} dp_j \right) + \sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} dH_i = 0, \quad (2.13)$$

then apply $N^* = P_0^{-1} P_f$ and substitute (2.13) in the resulting equation one gets

$$\begin{aligned} f_j \left(\frac{\partial \phi_j}{\partial q_j} dq_j + \frac{\partial \phi_j}{\partial p_j} dp_j \right) + \sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} N^* dH_i = \\ = -f_j \sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} dH_i + \sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} N^* dH_i = 0. \end{aligned}$$

It follows that

$$\sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} N^* dH_i = f_j \sum_{i=1}^n \frac{\partial \phi_j}{\partial H_i} dH_i, \quad j = 1, \dots, n,$$

that is, in matrix form,

$$N^* dH = dH F. \quad (2.14)$$

Here $n \times n$ control matrix F with eigenvalues $f_1, \dots, f_n$ is defined by

$$F = J^{-1} \text{diag}(f_1, \dots, f_n) J, \quad J_{ji} = \frac{\partial \phi_j}{\partial H_i}$$

and dH is $2n \times n$ matrix with entries $dH_{ij} = \partial H_j / \partial z_i$, where $z = (q, p)$.

Equation (2.14) means that the subspace spanned by covectors $dH_1, \dots, dH_n$ is invariant with respect to N^* [3].

Example 1 For further use we introduce the special control matrix F , which is the Fröbenius matrix

$$F_f = \begin{pmatrix} c_1 & 1 & 0 & \cdots & 0 \\ c_2 & 0 & 1 & \cdots & 0 \\ \vdots & & & & \\ c_n & 0 & \cdots & & 0 \end{pmatrix}. \quad (2.15)$$

Here c_k are coefficients of the characteristic polynomial $\Delta_N(\lambda)$ of the recursion operator $N = P_f P_0^{-1}$:

$$\Delta_N(\lambda) = \left(\det(N - \lambda I) \right)^{1/2} = \lambda^n - (c_1 \lambda^{n-1} + \dots + c_n) = \prod_{j=1}^n (\lambda - \lambda_j). \quad (2.16)$$

3 Open Toda lattice

Let us consider open Toda associated with the root system of $\mathcal{A}_n$ type. The Hamilton function is equal to

$$H = \frac{1}{2} \sum_{i=1}^n p_i^2 + \sum_{i=1}^{n-1} e^{q_i - q_{i+1}},$$

where q_i denotes the position of i -th particle and p_i is its momenta, such that

$$\{q_i, p_j\}_0 = \delta_{ij}, \quad \{p_i, p_j\}_0 = \{q_i, q_j\}_0 = 0. \quad (3.17)$$

Consequently, the equations of motion read

$$\dot{q}_i = p_i, \quad \dot{p}_1 = -e^{q_1 - q_2}, \quad \dot{p}_n = e^{q_{n-1} - q_n}$$

and

$$\dot{p}_i = e^{q_{i-1} - q_i} - e^{q_i - q_{i+1}}, \quad i = 2, \dots, n-1.$$

The exact solution is due to existence of a Lax matrix. Consider the L -operator

$$L_i = \begin{pmatrix} \lambda - p_i & -e^{q_i} \\ e^{-q_i} & 0 \end{pmatrix}, \quad (3.18)$$

and the monodromy matrix

$$T(\lambda) = L_1(\lambda) \cdots L_{n-1}(\lambda) L_n(\lambda) = \begin{pmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & D(\lambda) \end{pmatrix}, \quad \det T(\lambda) = 1, \quad (3.19)$$

which depends polynomially on the parameter λ

$$T(\lambda) = \begin{pmatrix} \lambda^n + A_1\lambda^{n-1} + \dots + A_n & B_1\lambda^{n-1} + \dots + B_n \\ C_1\lambda^{n-1} + \dots + C_n & D_1\lambda^{n-2} + \dots + D_n \end{pmatrix}.$$

The monodromy matrix satisfies Sklyanin's Poisson brackets:

$$\{T(\lambda) \otimes T(\mu)\}_0 = [r(\lambda - \mu), T(\lambda) \otimes T(\mu)], \quad (3.20)$$

where $r(\lambda - \mu)$ is the 4×4 rational r -matrix

$$r(\lambda - \mu) = \frac{-1}{\lambda - \mu} \Pi, \quad \Pi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \eta \in \mathbb{C}. \quad (3.21)$$

Monodromy matrix $T(\lambda)$ is the Lax matrix for periodic Toda lattice, whereas the Lax matrix for open Toda lattice is equal to

$$T_o(\lambda) = KT(\lambda) = \begin{pmatrix} A & B \\ 0 & 0 \end{pmatrix}(\lambda), \quad K = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

The trace of the Lax matrix

$$\text{tr } T_o(\lambda) = A(\lambda) = \lambda^n + H_1\lambda^{n-1} + \dots + H_n, \quad \{H_i, H_j\} = 0. \quad (3.22)$$

generates n independent integrals of motion H_i in the involution providing complete integrability of the system [7].

3.1 The Moser variables

According to [6] we introduce the n pairs of separated variables $\lambda_i, \mu_i, i = 1, \dots, n$, having the standard Poisson brackets,

$$\{\lambda_i, \lambda_j\}_0 = \{\mu_i, \mu_j\}_0 = 0, \quad \{\lambda_i, \mu_j\}_0 = \delta_{ij}, \quad (3.23)$$

with the λ -variables being n zeros of the polynomial $A(\lambda)$ and the μ -variables being values of the polynomial $B(\lambda)$ at those zeros,

$$A(\lambda_i) = 0, \quad \mu_i = \eta^{-1} \ln B(\lambda_i), \quad i = 1, \dots, n. \quad (3.24)$$

The Moser variables λ_j may be identified with poles of the Baker-Akhiezer function $\vec{\Psi}$ associated with the Lax matrix $T(\lambda)$ (3.19)

$$T(\lambda)\vec{\Psi} = \lambda\vec{\Psi}, \quad (\vec{\Psi}, \vec{\alpha}) = \text{const},$$

having some very special normalization $\vec{\alpha}$ [8].

The interpolation data (3.24) plus n identities

$$B(\lambda_i)C(\lambda_i) = \det T(\lambda) = 1$$

allow us to construct the separation representation for the whole monodromy matrix $T(\lambda)$:

$$\begin{aligned}
A(\lambda) &= (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n), \\
B(\lambda) &= A(\lambda) \sum_{i=1}^n \frac{e^{\mu_i}}{(\lambda - \lambda_i)A'(\lambda_i)}, \\
C(\lambda) &= -A(\lambda) \sum_{i=1}^n \frac{e^{-\mu_i}}{(\lambda - \lambda_i)A'(\lambda_i)}, \\
D(\lambda) &= \frac{1 + B(\lambda)C(\lambda)}{A(\lambda)}.
\end{aligned} \tag{3.25}$$

If we postulate the following second Poisson brackets (2.12)

$$\{\lambda_i, \mu_j\}_1 = \lambda_i \delta_{ij}, \quad \{\lambda_i, \lambda_j\}_1 = \{\mu_i, \mu_j\}_1 = 0$$

one gets [9]

$$\begin{aligned}
\{A(\lambda), A(\mu)\}_1 &= \{B(\lambda), B(\mu)\}_1 = 0, \\
\{A(\lambda), B(\mu)\}_1 &= \frac{1}{\lambda - \mu} \left(\mu A(\lambda)B(\mu) - \lambda A(\mu)B(\lambda) \right).
\end{aligned} \tag{3.26}$$

The first bracket in (3.26) guaranties that integrals of motion H_i (3.22) from $A(\lambda)$ are in the bi-involution.

Substitute polynomials $A(\lambda)$ and $B(\lambda)$ in initial (p, q) -variables into the brackets (3.26) and solve the resulting equations to obtain the known second Poisson tensor [1, 4]

$$P_1^{open} = \sum_{i=1}^{n-1} e^{q_i - q_{i+1}} \frac{\partial}{\partial p_{i+1}} \wedge \frac{\partial}{\partial p_i} + \sum_{i=1}^n p_i \frac{\partial}{\partial q_i} \wedge \frac{\partial}{\partial p_i} + \sum_{i < j}^n \frac{\partial}{\partial q_j} \wedge \frac{\partial}{\partial q_i}. \tag{3.27}$$

The minimal characteristic polynomial of the corresponding recursion operator $N_M = P_1^{open} P_0^{-1}$ is equal to

$$\Delta_{N_M}(\lambda) = A(\lambda) = \prod_{j=1}^n (\lambda - \lambda_j) = \lambda^n + \sum_{j=1}^n H_j \lambda^{n-j}.$$

So, (λ, μ) -coordinates are variables of separation of the action-angle type [3], i.e. the corresponding separated equations are trivial

$$\{H_i, \lambda_j\} = 0, \quad i, j = 1, \dots, n.$$

The Hamiltonians H_i (3.22) from $A(\lambda)$ satisfy the Fröbenius recursion relations

$$N_M^* dH_i = dH_{i+1} - H_i dH_1, \tag{3.28}$$

where $N_M^* = P_0^{-1} P_1^{open}$ and $H_{n+1} = 0$, i.e. Hamiltonians H_i satisfy equation (2.14) with the Fröbenius matrix F_f (2.15).

Example 2 For the 3-particles open Toda lattice Hamiltonians H_i (3.22) are

$$\begin{aligned} H_1 &= -(p_1 + p_2 + p_3), \\ H_2 &= p_1 p_2 + p_1 p_3 + p_2 p_3 - e^{q_1 - q_2} - e^{q_2 - q_3}, \\ H_3 &= -p_1 p_2 p_3 + p_1 e^{q_2 - q_3} + p_3 e^{q_1 - q_2}. \end{aligned}$$

It's obvious that $H = H_1^2/2 - H_2$. The second Poisson tensor P_1 (3.27) in the matrix form reads as

$$P_1^{open} = \begin{pmatrix} 0 & -1 & -1 & p_1 & 0 & 0 \\ 1 & 0 & -1 & 0 & p_2 & 0 \\ 1 & 1 & 0 & 0 & 0 & p_3 \\ -p_1 & 0 & 0 & 0 & -e^{q_1 - q_2} & 0 \\ 0 & -p_2 & 0 & e^{q_1 - q_2} & 0 & -e^{q_2 - q_3} \\ 0 & 0 & -p_3 & 0 & e^{q_2 - q_3} & 0 \end{pmatrix}. \quad (3.29)$$

The control matrix F in (2.14) is the Fröbenius matrix

$$F_M^{open} = \begin{pmatrix} -H_1 & 1 & 0 \\ -H_2 & 0 & 1 \\ -H_3 & 0 & 0 \end{pmatrix}, \quad (3.30)$$

where coefficients $c_i = -H_i$ coincide with integrals of motion.

3.2 The Sklyanin variables

According to [7, 8] we can consider another set of the separated coordinates, which are poles of the Baker-Akhiezer function $\vec{\Psi}$ associated with the Lax matrix $T(\lambda)$ (3.19) having the standard normalization $\vec{\alpha} = (0, 1)$.

In this case the first half of variables is coming from $(n - 1)$ finite roots and logarithm of leading coefficient of the non-diagonal entry of the monodromy matrix

$$B(\lambda) = -e^{u_n} \prod_{j=1}^{n-1} (\lambda - v_j), \quad (3.31)$$

Another half is given by

$$u_j = -\ln A(v_j), \quad j = 1, \dots, n-1, \quad \text{and} \quad v_n = \sum_{i=1}^n p_i. \quad (3.32)$$

In these separated variables other entries of $T(\lambda)$ read as

$$A(\lambda) = \left(\lambda + \sum_{j=1}^n v_j \right) \prod_{j=1}^{n-1} (\lambda - v_j) + \sum_{j=1}^{n-1} e^{-u_j} \prod_{i \neq j}^{n-1} \frac{\lambda - v_i}{v_j - v_i}. \quad (3.33)$$

and

$$D(\lambda) = - \sum_{j=1}^{n-1} e^{u_j} \prod_{i \neq j}^{n-1} \frac{\lambda - v_i}{v_j - v_i}, \quad C(\lambda) = \frac{A(\lambda)D(\lambda) - 1}{B(\lambda)}.$$

If we postulate the second Poisson brackets (2.12)

$$\{v_i, u_j\}_1^* = v_i \delta_{ij}, \quad \{v_i, v_j\}_1^* = \{u_i, u_j\}_1^* = 0$$

one gets

$$\{A(\lambda), A(\mu)\}_1^* = \{B(\lambda), B(\mu)\}_1^* = 0, \quad (3.34)$$

and

$$\begin{aligned} \{A(\lambda), B(\mu)\}_1^* &= \frac{1}{\lambda - \mu} \left(\lambda A(\lambda) B(\mu) - \mu A(\mu) B(\lambda) \right) \\ &+ e^{-u_n} \left(\lambda + \mu + \sum_{i=1}^{n-1} v_i \right) B(\lambda) B(\mu). \end{aligned} \quad (3.35)$$

In initial (p, q) -variables the last bracket looks like

$$\begin{aligned} \{A(\lambda), B(\mu)\}_1^* &= \frac{1}{\lambda - \mu} \left(\lambda A(\lambda) B(\mu) - \mu A(\mu) B(\lambda) \right) \\ &+ e^{-q_n} \left(\lambda + \mu + \sum_{i=1}^{n-1} p_i \right) B(\lambda) B(\mu). \end{aligned} \quad (3.36)$$

The first bracket in (3.34) guaranties that integrals of motion H_i (3.22) from $A(\lambda)$ are in the bi-involution.

Substitute into the brackets (3.34)-(3.36) polynomials $A(\lambda)$ and $B(\lambda)$ in initial (p, q) -variables and solve the resulting equations to obtain the following Poisson tensor

$$\begin{aligned} P_1^* &= \sum_{i=1}^{n-2} e^{q_i - q_{i+1}} \frac{\partial}{\partial p_{i+1}} \wedge \frac{\partial}{\partial p_i} + \sum_{i=1}^{n-1} p_i \frac{\partial}{\partial q_i} \wedge \frac{\partial}{\partial p_i} + \sum_{i < j}^{n-1} \frac{\partial}{\partial q_j} \wedge \frac{\partial}{\partial q_i} \\ &+ \sum_{i=1}^{n-1} (p_i + p) \frac{\partial}{\partial p_n} \wedge \frac{\partial}{\partial q_i} + \sum_{i=2}^{n-1} (e^{q_i - q_{i+1}} - e^{q_{i-1} - q_i}) \frac{\partial}{\partial p_n} \wedge \frac{\partial}{\partial p_i} \\ &+ p \frac{\partial}{\partial p_n} \wedge \frac{\partial}{\partial q_n} + e^{q_1 - q_2} \frac{\partial}{\partial p_1} \wedge \frac{\partial}{\partial p_n} + e^{q_{n-2} - q_{n-1}} \frac{\partial}{\partial p_{n-1}} \wedge \frac{\partial}{\partial p_n}, \end{aligned} \quad (3.37)$$

where $p = \sum_{i=1}^n p_i$ is a total momentum.

The tensor P_1^* is independent on q_n and the minimal characteristic polynomial of the corresponding recursion operator $N_S = P_1^* P_0^{-1}$ is equal to

$$\Delta_{N_S} = -e^{q_n} (\lambda + p) B(\lambda).$$

The normalized traces of the powers of N_S are integrals of motion for $n - 1$ particle open Toda lattice. As consequence the Hamiltonians H_i (3.22) from $A(\lambda)$ satisfy equation (2.14) with the following control matrix

$$F_S^{open} = \begin{pmatrix} -p & 0 \\ 0 & F_M^{open} \end{pmatrix}, \quad (3.38)$$

where F_M^{open} is the Fröbenius matrix (2.15) associated with the recursion operator N_M for $n - 1$ particle open Toda lattice.

The corresponding separated equations follow directly from the definitions of (u, v) -variables (3.32)

$$e^{-u_j} - A(v_j) = 0, \quad j = 1, \dots, n - 1 \quad \text{and} \quad \Phi_n = v_n - \alpha_1 = 0, \quad (3.39)$$

where $A(\lambda) = \lambda^n + \sum_{i=1}^{n-1} \alpha_i \lambda^{n-i}$ and $\alpha_i = H_i$ are the values of integrals of motion. The first $(n - 1)$ separated equations give rise to the equations of motion

$$\{A(\lambda), v_j\} = A(v_j) \prod_{i \neq j}^{n-1} \frac{\lambda - v_i}{v_j - v_i}, \quad j = 1, \dots, n - 1, \quad (3.40)$$

which are linearized by the Abel transformation [7]

$$\left\{ A(\lambda), \sum_{k=1}^{n-1} \int^{v_k} \sigma_j \right\} = -\lambda^{j-1}, \quad \sigma_j = \frac{\lambda^{j-1} d\lambda}{A(\lambda)}, \quad j = 1, \dots, n - 1,$$

where $\{\sigma_j\}$ is a basis of abelian differentials of first order on an algebraic curve $z = A(\lambda)$ corresponding to the separated equations (3.39).

Remark 4 From the factorization of the monodromy matrix $T(\lambda)$ (3.19) one gets

$$T_n(\lambda) = T_{n-1}(\lambda) L_n(\lambda), \quad \Rightarrow \quad B_n(\lambda) = -e^{q_n} A_{n-1}(\lambda),$$

where $B_n(\lambda)$ is entry of the monodromy matrix $T_n(\lambda)$ of n particles Toda lattice and $A_{n-1}(\lambda)$ is entry of the monodromy matrix $T_{n-1}(\lambda)$ of $n - 1$ particles Toda lattice.

This implies that for the $(n - 1)$ -particle chain the Moser variables λ_j coincide with the $n - 1$ Sklyanin variables u_j , $i = 1, \dots, n - 1$ for the n -particle chain.

Example 3 At $n = 3$ the Poisson tensor P_1^* in the matrix form reads as

$$P_1^* = \begin{pmatrix} 0 & -1 & 0 & p_1 & 0 & -p_1 - p \\ 1 & 0 & 0 & 0 & p_2 & -p_2 - p \\ 0 & 0 & 0 & 0 & 0 & p \\ -p_1 & 0 & 0 & 0 & -e^{q_1 - q_2} & e^{q_1 - q_2} \\ 0 & -p_2 & 0 & e^{q_1 - q_2} & 0 & -e^{q_1 - q_2} \\ p_1 + p & p_2 + p & p & -e^{q_1 - q_2} & e^{q_1 - q_2} & 0 \end{pmatrix}.$$

and at $n=4$ it looks like

$$P_1^* = \begin{pmatrix} 0 & -1 & -1 & 0 & p_1 & 0 & 0 & -p_1 - p \\ 1 & 0 & -1 & 0 & 0 & p_2 & 0 & -p_2 - p \\ 1 & 1 & 0 & 0 & 0 & 0 & p_3 & -p_3 - p \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & p \\ -p_1 & 0 & 0 & 0 & 0 & -e^{q_1 - q_2} & 0 & e^{q_1 - q_2} \\ 0 & -p_2 & 0 & 0 & e^{q_1 - q_2} & 0 & -e^{q_2 - q_3} & e^{q_2 - q_3} - e^{q_1 - q_2} \\ 0 & 0 & -p_3 & 0 & 0 & e^{q_2 - q_3} & 0 & e^{q_2 - q_3} \\ p_1 + p & p_2 + p & p_3 + p & p & -e^{q_1 - q_2} & -e^{q_2 - q_3} + e^{q_1 - q_2} & -e^{q_2 - q_3} & 0 \end{pmatrix}.$$

The corresponding control matrices F in (2.14) are given by

$$F_S^{open} = \begin{pmatrix} -p_1 - p_2 - p_3 & 0 & 0 \\ 0 & p_1 + p_2 & 1 \\ 0 & -p_1 p_2 + e^{q_1 - q_2} & 0 \end{pmatrix}. \quad (3.41)$$

and

$$F_S^{open} = \begin{pmatrix} -p_1 - p_2 - p_3 - p_4 & 0 & 0 & 0 \\ 0 & p_1 + p_2 + p_3 & 1 & 0 \\ 0 & -p_1 p_2 - p_1 p_3 - p_2 p_3 + e^{q_1 - q_2} - e^{q_2 - q_3} & 0 & 1 \\ 0 & p_1 p_2 p_3 - p_1 e^{q_2 - q_3} - p_3 e^{q_1 - q_2} & 0 & 0 \end{pmatrix}. \quad (3.42)$$

3.3 Periodic Toda lattice

The trace of the Lax matrix $T(\lambda)$ (3.19) for periodic Toda lattice

$$\text{tr } T(\lambda) = A(\lambda) + D(\lambda) = \lambda^n + H_1 \lambda^{n-1} + \cdots H_n, \quad \{H_i, H_j\} = 0. \quad (3.43)$$

generates n independent integrals of motion H_i in the involution providing complete integrability of the periodic Toda lattice [7]. For instance the Hamilton function is equal to

$$H = \frac{1}{2} \sum_{i=1}^n p_i^2 + \sum_{i=1}^{n-1} e^{q_i - q_{i+1}} + e^{q_n - q_1}.$$

The Moser (λ, μ) -variables (3.24) are the separated variables for the open Toda lattice only. Nevertheless, the bi-Hamiltonian structure for the periodic Toda lattice may be constructed by using generic properties of the Sklyanin bracket (3.20) [10]. According to [10], in order to get the second Poisson tensor P_1^{per} for the periodic Toda lattice we have to apply canonical transformation

$$p_1 \rightarrow p_1 + e^{-q_1}, \quad p_n \rightarrow p_n + e^{q_n}. \quad (3.44)$$

to the tensor P_1^{open} (3.27) for the open Toda lattice. In this case Hamiltonians (3.43) satisfy equation (2.14) with the standard Fröbenius matrix F_f (2.15) associated with the recursion operator $N_M^{per} = P_1^{per} P_0^{-1}$ [10].

The Sklyanin (u, v) -variables (3.31)-(3.32) are the separated variables for the open and periodic Toda lattice simultaneously. Therefore, the one Poisson tensor P_1^* determines the bi-Hamiltonian structure for the both Toda lattices, but the corresponding Hamiltonians satisfy equation (2.14) with the slightly different control matrices F_S^{open} and F_S^{per} .

Namely, for the open Toda lattice the control matrix F_S^{open} is given by (3.38). For the periodic Toda lattice we have to change the first column of this control matrix F_S^{open} (3.38) by the rule:

$$F_{1,1}^{per} = -p, \quad F_{i,1}^{per} = (p + \sum_{j=1}^{n-1} p_j) F_{i,2}^{open} + F_{i+1,2}^{open}, \quad i = 2, \dots, n, \quad (3.45)$$

where $F_{n+1,2}^{open} = 0$.

Remark 5 For the open and periodic Toda lattices we have two matrix equations for the same integrals of motion:

$$N_M^* dH = dH F_M, \quad N_S^* dH = dH F_S$$

So, we have various complimentary relations

$$N_M^* N_S^* dH = dH F_M F_S, \quad \text{or} \quad N_S^* (N_M^*)^{-1} dH F_M = dH F_S.$$

between integrals of motion H_i , recursion operators N_S , N_M and the corresponding control matrices F_M , F_S .

Example 4 For the 3-particles periodic Toda lattice Hamiltonians H_i (3.43) are

$$\begin{aligned} H_1 &= -(p_1 + p_2 + p_3), \\ H_2 &= p_1 p_2 + p_1 p_3 + p_2 p_3 - e^{q_1 - q_2} - e^{q_2 - q_3} - e^{q_3 - q_1}, \\ H_3 &= -p_1 p_2 p_3 + p_1 e^{q_2 - q_3} + p_2 e^{q_3 - q_1} + p_3 e^{q_1 - q_2}. \end{aligned}$$

The second Poisson tensor (3.29) after canonical transformation (3.44) reads as

$$P_1^{per} = \begin{pmatrix} 0 & -1 & -1 & e^{-q_1} + p_1 & 0 & e^{q_3} \\ 1 & 0 & -1 & e^{-q_1} & p_2 & e^{q_3} \\ 1 & 1 & 0 & e^{-q_1} & 0 & p_3 + e^{q_3} \\ -e^{-q_1} - p_1 & -e^{-q_1} & -e^{-q_1} & 0 & -e^{q_1 - q_2} & e^{q_3 - q_1} \\ 0 & -p_2 & 0 & e^{q_1 - q_2} & 0 & -e^{q_2 - q_3} \\ -e^{q_3} & -e^{q_3} & -p_3 - e^{q_3} & -e^{q_3 - q_1} & e^{q_2 - q_3} & 0 \end{pmatrix}.$$

The corresponding control matrix F in (2.14) is the Fröbenius matrix

$$F_M^{per} = \begin{pmatrix} c_1 & 1 & 0 \\ c_2 & 0 & 1 \\ c_3 & 0 & 0 \end{pmatrix}, \quad (3.46)$$

where

$$c_1 = p_1 + e^{-q_1} + p_2 + p_3 + e^{q_3},$$

$$c_2 = -p_1 p_2 - p_1 p_3 - p_2 p_3 - (p_2 + p_3)e^{-q_1} - (p_1 + p_2)e^{q_3} + e^{q_1 - q_2} + e^{q_2 - q_3} - e^{q_1 - q_3},$$

$$c_3 = ((p_3 + e^{q_3})p_2 - e^{q_2 - q_3})p_1 + (p_3 + e^{q_3})p_2 e^{-q_1} - p_3 e^{q_1 - q_2} - e^{-q_1 + q_2 - q_3} - e^{q_1 - q_2 + q_3}.$$

are coefficients of the minimal characteristic polynomial of the corresponding recursion operator $N_M^{per} = P_1^{per} P_0^{-1}$.

The same integrals of motion H_i for periodic Toda lattice satisfy equation (2.14) with another recursion operator $N_S^* = P_0^{-1} P_1^*$ and with the following control matrix (3.38)-(3.45)

$$F_S^{per} = \begin{pmatrix} -p_1 - p_2 - p_3 & 0 & 0 \\ e^{q_1 - q_2} - p_1 p_2 + (p_1 + p_2)(2p_1 + 2p_2 + p_3) & p_1 + p_2 & 1 \\ (e^{q_1 - q_2} - p_1 p_2)(2p_1 + 2p_2 + p_3) & e^{q_1 - q_2} - p_1 p_2 & 0 \end{pmatrix}. \quad (3.47)$$

4 Conclusion.

For the Toda lattice associated with the root system of $\mathcal{A}_n$ type we present two different bi-Hamiltonian structures on $M \simeq \mathbb{R}^{2n}$. The introduced Poisson tensors P_1 (3.27) and P_1^* (3.37) are incompatible

$$[[P_1, P_1^*]] \neq 0,$$

while the corresponding recursion operators $N_M = P_1 P_0^{-1}$ and $N_S = P_1^* P_0^{-1}$ take the diagonal form in the Moser and the Sklyanin separated variables respectively.

Associated with the tensor P_1 brackets (3.26) were rewritten in the r -matrix form in [10]. It will be interesting to get the similar r -matrix formulation for the brackets (3.34)-(3.36) associated with the tensor P_1^* .

The research was partially supported by the RFBR grant 06-01-00140 and grant NSc5403.2006.1.

References

- [1] A. Das, S. Okubo, *A systematic study of the Toda lattice*, Ann. Phys., v.30, p.215-232, 1989.
- [2] G. Falqui, F. Magri, M. Pedroni, *Bihamiltonian geometry and separation of variables for Toda lattices*, J. Nonlinear Math. Phys., v.8, p.118-127, 2001.
- [3] G. Falqui, M. Pedroni, *Separation of variables for bi-Hamiltonian systems*, Math. Phys. Anal. Geom., v.6, p.139-179, 2003.
- [4] R. L. Fernandes, *On the master symmetries and bi-Hamiltonian structure of the Toda lattice*, J. Phys. A, v.26, p.3797-3803, 1993.
- [5] F. Magri, *Eight lectures on Integrable Systems*, Lecture Notes in Physics, Springer Verlag, Berlin-Heidelberg, v.495, p.256-296, 1997.
- [6] J. Moser, *Finitely many mass points on the line under the influence of an exponential potential - an integrable system*, Lect. Notes in Phys., v.38, p.467-497, 1975.
- [7] E.K. Sklyanin, *The quantum Toda chain*, Lect. Notes in Physics, v.226, p.196-293, 1985.
- [8] E.K. Sklyanin, *Separation of variables—new trends*, Progr. Theor. Phys. Suppl., v.118, p.35, 1995.
- [9] A. V. Tsiganov, Yu.A. Grigoryev, *On the Darboux-Nijenhuis variables for the open Toda lattice*, in the Vadim Kuznetsov Memorial Issue of SIGMA, v.2, paper 097, nlin.SI/0701004, 2006.
- [10] A.V. Tsiganov *A family of the Poisson brackets compatible with the Sklyanin bracket*, submitted to J. Phys.A, nlin.SI/0612025, 2006.